

FTD Module

- We would like to ask you a number of questions about changes in the patient's behaviour that you may have noticed. Your view is important as it will help us in our assessment. Please read each question carefully. For each question, indicate the *frequency* of the behaviour, followed by the level of *severity*.
- All questions apply to the patient's behaviour over the past one month.**

	Does this symptom occur?	How often?				How severe?		
		Less than weekly	About once per week	Several times a week	Daily or continuously	Mild	Moderate	Severe
		1	2	3	4	1	2	3
Loss of sympathy/empathy								
Does the patient seem to respond less to other people's needs or feelings? Is there reduced personal warmth? Do they seem to lack empathy or sympathy for others?	Yes <input type="checkbox"/> (then answer how often and how severe) No <input type="checkbox"/> (then proceed to next question)	Less than weekly	About once per week	Several times a week	Daily or continuously	Loss of sympathy or empathy is notable but is only mildly different to patient's usual behaviour	Loss of sympathy or empathy is very evident and occurs in a substantial number of interactions with people	Loss of sympathy or empathy is very evident and occurs in the majority of interactions with other people
Ritualistic/compulsive behaviour								
Does the patient engage in compulsive behaviours such as continually checking, counting or cleaning things? Does the patient have fixed rituals or routines that they must perform?	Yes <input type="checkbox"/> (then answer how often and how severe) No <input type="checkbox"/> (then proceed to next question)	Less than weekly	About once per week	Several times a week	Daily or continuously	Ritualistic or compulsive behaviours are notably present but are not disturbing	Ritualistic or compulsive behaviours are very evident but may be able to be overcome by the caregiver	Ritualistic or compulsive behaviours are very evident and usually do not respond to any intervention by the caregiver
Poor response to social/emotional cues								
Does the patient seem to be worse at interacting socially with others e.g. is there less eye contact or do they stand closer than normal to others? Are they insensitive to how a normal conversation flows? Do they seem to have difficulty reading other's emotions?	Yes <input type="checkbox"/> (then answer how often and how severe) No <input type="checkbox"/> (then proceed to next question)	Less than weekly	About once per week	Several times a week	Daily or continuously	Pays noticeably less attention to social cues, or sometimes responds awkwardly or unexpectedly to social cues	Pays much less attention to social cues, or often responds awkwardly or unexpectedly to social cues	Pays almost no attention to social cues, or very often responds awkwardly or unexpectedly to social cues
Inappropriate trusting behaviour								
Does the patient seem more gullible e.g. have they been taken advantage of? Are they less cautious than before? Have they performed any acts of poor judgment e.g. fallen for scams or purchased something extravagant they wouldn't have done before?	Yes <input type="checkbox"/> (then answer how often and how severe) No <input type="checkbox"/> (then proceed to next question)	Less than weekly	About once per week	Several times a week	Daily or continuously	Has displayed a few clear but minor acts of poor judgment of other people	Has displayed multiple minor acts or a few major acts of poor judgment of other people resulting in adverse consequences	Has displayed recurrent, in at least some cases severe, acts of poor judgment of other people resulting in adverse consequences

Hyperreligiosity								
Does the patient seem to have developed increased or unusual religious experiences or feelings compared to previously?	Yes <input type="checkbox"/> (then answer how often and how severe) No <input type="checkbox"/> (then proceed to next question)	Less than weekly	About once per week	Several times a week	Daily or continuously	Increased or unusual religious experiences are present but seem harmless and produce little distress	Increased or unusual religious experiences may be distressing and disruptive	Increased or unusual religious experiences are very disruptive and are a major source of behavioural disruption
Hypersexuality								
Has the patient started to engage in unusual or excessive sexual behaviour? Do they have increased libido compared to before?	Yes <input type="checkbox"/> (then answer how often and how severe) No <input type="checkbox"/> (then proceed to next question)	Less than weekly	About once per week	Several times a week	Daily or continuously	Sexual behaviour is disruptive but can be managed by redirection or reassurance	Sexual behaviour is disruptive and difficult to redirect or control	Sexual behaviour is very disruptive and a major source of difficulty
Altered sense of humour								
Has the patient developed a change in sense of humour, such as finding things funny which they didn't before or finding things funny that others don't?	Yes <input type="checkbox"/> (then answer how often and how severe) No <input type="checkbox"/> (then proceed to next question)	Less than weekly	About once per week	Several times a week	Daily or continuously	Mild but definite change in sense of humour. May find things funny which others don't	Sense of humour completely different from previously (e.g. may prefer more physical or slapstick humour)	Complete loss of sense of humour
Altered responsiveness to pain and/or temperature								
Has the way that the patient seems to perceive pain or temperature changed? Do they seem to have a reduced or heightened response to pain? Do they seem to have changed in how they tolerate heat or cold (e.g. wearing less clothes when it's cold, or more clothes when it's hot)?	Yes <input type="checkbox"/> (then answer how often and how severe) No <input type="checkbox"/> (then proceed to next question)	Less than weekly	About once per week	Several times a week	Daily or continuously	Mild change in responsiveness to pain or temperature – no limitation on daily activities	Moderate change in responsiveness to pain or temperature – some limitation on daily activities	Severe change in responsiveness to pain or temperature – limits most daily activities